

# Treatment Efficacy of Benign Paroxysmal Positional Vertigo (BPPV) with Canalith Repositioning Maneuver and Semont Liberatory Maneuver in 376 Patients

Richard E. Gans, Ph.D., FAAA,<sup>1</sup> and  
Patricia A. Harrington-Gans, Au.D., FAAA<sup>1</sup>

## ABSTRACT

---

Canalith Repositioning and Semont Liberatory Maneuvers have been shown to be highly efficacious in the successful treatment of Benign Paroxysmal Positional Vertigo (BPPV). The differentiation of canalolithiasis, cupulolithiasis, and correct identification of canal involvement, particularly through the use of Video-oculography, have enhanced treatment decisions and outcomes.

Since 1994, approximately 700 BPPV patients have been treated at the authors' clinical facility. An anecdotal study of 376 of these patients followed over a 7-year period is presented. The patients in the historical study all presented with BPPV- PC and were treated with modified Canalith Repositioning Maneuver and Semont Liberatory Maneuver treatment procedures. The review indicated no significant differences in treatment outcomes between the two procedures. Seventy-nine percent of the patients required only one treatment, while 17% required two treatments, 3.5% required three treatments, and 0.05% required four treatments. The average number of treatments was 1.3. The SLM did show a reduced recurrence rate compared to the CRP method.

**KEYWORDS:** Benign Paroxysmal Positional Vertigo (BPPV), canalolithiasis, cupulolithiasis, otolith

**Learning Outcomes:** As a result of this activity, the reader will: (1) have a historical perspective of the development of the diagnosis and treatment of BPPV; (2) have a review of the literature of BPPV treatment methodologies; and (3) be able to determine and perform appropriate BPPV treatment methods.

---

Vestibular Diagnosis and Rehabilitation: Science and Clinical Applications; Editor in Chief, Catherine V. Palmer, Ph.D.; Guest Editor, Richard E. Gans, Ph.D., FAAA. *Seminars in Hearing*, volume 23, number 2, 2002. Address for correspondence and reprint requests: Richard E. Gans, Ph.D., The American Institute of Balance, 11290 Park Boulevard, Seminole, FL 33772. E-mail: rgans@dizzy.com. <sup>1</sup>The American Institute of Balance, Seminole, Florida. Copyright © 2002 by Thieme Medical Publishers, Inc., 333 Seventh Avenue, New York, NY 10001, USA. Tel: +1(212) 584-4662. 00734-0451,p;2002,23,02,129,142,ftx,en;sih00204x.

Benign paroxysmal positional vertigo (BPPV) is the most common form of vertigo<sup>1</sup> and is among the most easily treated without the need for surgery or medication.<sup>2</sup> The past 14 years has provided clinicians a wealth of research, retrospective studies, identification of BPPV variants and new treatment techniques to consider and practice. This article will present a review of, as well as the author's experience with, the treatment efficacy for posterior canal BPPV with both the Semont Liberatory and Canalith Repositioning Maneuvers in treating 376 patients over an 8-year period. A historical perspective will precede a comprehensive discussion of clinical protocols, and a review and discussion of the anecdotal findings.

## DESCRIPTION AND PATHOPHYSIOLOGY

It has been 80 years since Barany<sup>3</sup> first described the abnormal condition eliciting rotatory-torsional nystagmus and vertigo with change in head position, a condition we now know as BPPV. It would be 30 years before Dix and Hallpike would name it Benign Paroxysmal Positional Vertigo (BPPV) in 1952.<sup>4</sup> Their work contributed to our understanding of the positioning maneuver to elicit the response and the cardinal manifestations. It did not explain its underlying pathophysiology. There was much controversy as to the cause of the condition until Schuknecht's classic cupulolithiasis explanation was published in 1969.<sup>5</sup> Prior to that time, many thought that the rotatory-torsional nystagmus and transient intense vertigo was too complicated to be attributable to anything but a central nervous system disorder.<sup>6</sup>

Schuknecht's histological and temporal bone studies demonstrated that the otoconia from the utricle migrated into the posterior semicircular canal (PSCC) and embedded onto the cupula. The weighted cupula, when moved from the vertical to the horizontal plane, which occurs when the head is tilted back, would cause the posterior canal to inadvertently become a gravity sensor. Although Schuknecht's work advanced the theory of the biomechanical aspects of BPPV, it did not completely ex-

plain all aspects of the condition. Ten years later, the term canalithiasis was coined by Parnes and McClure.<sup>7</sup> While attempting to blockade the ampullae of the PSCC, they observed the presence of free-floating otoconia within the long process of the PSCC. The biomechanics of this form of BPPV readily explains the fatigability aspect of the response, as had previously been postulated by Hall et al.<sup>8</sup> It is now accepted that while BPPV may manifest itself in a variety of variants and/or combinations, canalithiasis is the most prevalent form of BPPV.<sup>9</sup>

## THERAPEUTIC TREATMENT METHODS

A physical therapeutic maneuver to specifically ameliorate BPPV was first proposed by Brandt and Daroff in 1980.<sup>10</sup> Their repetitive side-lying maneuver, although less than completely efficacious, did have some degree of success in reducing or eliminating the BPPV response in some patients. The self-induced vertigo, and uncomfortable and unnatural positions of the patient, limits its efficacy. Various techniques and successes in the treatment of BPPV have been put forth, beginning with the Liberatory maneuver by Semont, Freyss and Vitte<sup>11</sup> in 1988, and followed by the Canalith Repositioning by Epley.<sup>12</sup> The reported success in 1993 of Herdman et al<sup>13</sup> using the single treatment technique began a litany of reports of successful treatment options by more than a score of researchers. Table 1<sup>11-24,27,29,39,44</sup> provides a review of these studies.

Most clinicians who perform Repositioning and Liberatory Maneuvers have modified many of the protocols into a hybrid that has produced successful outcomes for them in their own clinics. It appears that regardless of what methodology is used, if the differential diagnosis of canalithiasis from cupulolithiasis is made, and appropriate canal involvement has been obtained, there is high treatment efficacy.<sup>14-24</sup>

Treatment of BPPV by either Liberatory or Canalith Repositioning Maneuver is widely recognized as the most efficacious, noninvasive and cost effective way to manage the condi-

**Table 1 Studies of Posterior Canal Benign Paroxysmal Positional Vertigo Treatment Methods and Efficacy**

Study	N	Year	Procedure	Results
Norre and Beckers <sup>45</sup>	51	1988	Other	Vestibular habituation training produces good results, when Benign Paroxysmal Positional Vertigo is diagnosed correctly.
Semont et al <sup>11</sup>	711	1988	Semont Liberatory	Semont Liberatory Maneuver is an effective form of treatment showing a low recurrence rate.
Epley <sup>12</sup>	30	1992	Canalith Repositioning Maneuver	Canalith Repositioning Maneuver, as described by Epley, is the most effective form of treatment.
Troost & Patton <sup>46</sup>	N/A	1992	Other	There is an excellent chance that exercise therapy will be curative with patients having Benign Paroxysmal Positional Vertigo.
Herdman et al <sup>13</sup>	60	1993	Semont Liberatory and Canalith Repositioning Maneuvers	Both the Semont and Epley maneuvers are effective treatments for Benign Paroxysmal Positional Vertigo.
Steenerson and Cronin <sup>14</sup>	40	1996	Canalith Repositioning	Canalith Repositioning Maneuver is as effective as vestibular habituation training and requires less patient time for treatment.
Coppo et al <sup>15</sup>	165	1996	Semont Liberatory Maneuver	Over 80% of patients were successfully treated after 1 to 3 sessions.
Herdman & Tusa <sup>29</sup>	85	1996	Canalith Repositioning Maneuver	Canalith Repositioning Maneuver can cause debris to move into the anterior or horizontal canal.
Lempert et al <sup>16</sup>	15	1997	Other	360° rotation of the posterior/semicircular canal proved to be an effective treatment.
Cohen and Jerabek <sup>17</sup>	87	1999	Both	Found that augmented head rotations are unnecessary and the modified Epley and Semont maneuvers are equally effective in the treatment of benign paroxysmal positional vertigo.
Gall et al <sup>18</sup>	16	1999	Semont Liberatory Maneuver	The findings showed a statistically significant change in subjective visual vertical post Hallpike and Semont maneuvers.

**Table 1 (Continued)**

<b>Study</b>	<b>N</b>	<b>Year</b>	<b>Procedure</b>	<b>Results</b>
Wolf et al <sup>19</sup>	107	1999	Canalith Repositioning Maneuver	The modified Epley maneuver is an excellent treatment for benign paroxysmal positional vertigo.
Blatt et al <sup>47</sup>	33	2000	Canalith Repositioning Maneuver	Treatment with Canalith Repositioning Maneuver appears to improve balance, but not in all patients.
Dornhoffer and Colvin <sup>20</sup>	52	2000	Canalith Repositioning Maneuver	Canalith Repositioning Maneuver was effective in 99% of patients. Any recurrence is thought to correlate to etiology.
Gross et al <sup>48</sup>	9	2000	Canalith Repositioning Maneuver	Meniere's disease may predispose patients to intractable benign paroxysmal positional vertigo.
Macias et al <sup>21</sup>	259	2000	Canalith Repositioning Maneuver	Patients with benign paroxysmal positional vertigo not located in a single posterior semicircular canal are more likely to need multiple Canalith Repositioning procedures.
Nuti et al <sup>33</sup>	56	2000	Semont Liberatory Maneuver	Restrictions following Semont Liberatory Maneuver are not necessary when treating benign paroxysmal positional vertigo.
Nunez et al <sup>22</sup>	168	2000	Canalith Repositioning Maneuver	Canalith Repositioning Maneuver is an effective treatment for benign paroxysmal positional vertigo with only 1 or 2 sessions; however, there is a 15% recurrence of benign paroxysmal positional vertigo symptoms per year after treatment.
O'Reilly et al <sup>23</sup>	72	2000	Canalith Repositioning Maneuver	Benign paroxysmal positioning vertigo is effective in relieving the vertigo associated with benign paroxysmal positional vertigo.
Tirelli et al <sup>24</sup>	118	2000	Canalith Repositioning Maneuver	Success rates were significantly higher with the Modified Repositioning Maneuver.
Sargent et al <sup>30</sup>	168	2001	Canalith Repositioning Maneuver and other	Mastoid oscillation does not significantly improve the efficacy of the Canalith Repositioning Maneuver.

tion.<sup>2</sup> The high incidence of BPPV among the elderly persons has been documented. An incidence rate as great as 50% in persons over the age of 70, and the associated morbidity with undiagnosed and untreated BPPV has been established.<sup>25</sup> Gans and Crandell<sup>26</sup> have reported significant improvement in the subjective report (SF-36)<sup>27</sup> of BPPV patients post-treatment in quality of life, general health, mental health, and vitality.

### TECHNICAL ADVANCES

Recent advances in technology, specifically infrared video-oculography, have provided clinicians with the capability to record and review eye movements associated with BPPV. This has allowed better differentiation of the involved canal(s). Approximately 90 to 95% of BPPV will affect the PSCC and, to a lesser degree, the HSCC.<sup>28</sup> The predominant occurrence of HC-BPPV is usually secondary to the treatment of PC-BPPV with migration due to the repositioning itself. The occurrence of AC-BPPV rarely is seen or reported. Its existence may be questionable in view of the anatomical position of the anterior canal. When it does occur, it may appear as a cupulolithiasis variant.

The use of this new technology gives the clinician the opportunity of repeated viewing of the associated eye movement. The otherwise brief and transitory nature of the clinician's observation of the nystagmus in real time may be insufficient or inaccurate for a differential diagnosis. Figure 1 is a screen display of a computerized videonystagmography (VNG) system. A differential diagnostic clinical pathway is presented in Figure 2 for canalolithiasis vs. cupulolithiasis.

### TREATMENT OF CANALITHIASIS

Canalithiasis may be treated using either the Canalith Repositioning Maneuver (Appendix A), or the Semont Liberatory Maneuver (Appendix B). The decision of which method to follow becomes a matter of comfort or choice

for the clinician. The patient's physical characteristics or capabilities need to be taken into consideration. The Liberatory Maneuver requires the movement of the patient en masse and may be inappropriate for those patients who have undergone a hip replacement within 90 days of the treatment. The Canalith Repositioning Maneuver tends to be more comfortable for patients, as it only requires them to move their head and roll onto their side. Physical limitations, whether they are orthopedic, neuromuscular, or due to obesity, may make one technique easier and more comfortable for both patient and clinician than the other. The success rate of either procedure has been reported to be greater than 90% by most investigators.

### TREATMENT OF CUPULOLITHIASIS

According to Schuknechts<sup>10</sup> theory of cupulolithiasis, the debris adheres to the cupula rather than free floating in the long process of the posterior canal. This theory, according to the Semont's Liberatory Maneuver, recognizes that the debris cannot merely be repositioned by rotation of the head but some basic principals of physics must be used to dislodge the debris from the cupula so that it can then be released and allowed to return to the utricle and dissolve. The differentiating factor and the diagnosis are based on whether the symptoms will abate with repetition of a provoking maneuver. For patients with acute symptoms, including nausea and emesis, repetition of three and four modified Hallpike Maneuvers as part of the differential diagnostic process may be difficult.

### USE OF VIBRATION

The original work of Epley,<sup>12</sup> as well as Li,<sup>29</sup> recommends the use of vibration to maximize treatment outcome. However, a review of the literature suggests that most researchers and clinicians have not found the vibrator to be a critical component in the treatment of BPPV.<sup>13-24</sup> There are other options that range from bone conduction oscillators, tuning forks,





Figure 1 Videoculography technology allows visualization and video recording of eye movement in addition to the computer graphics.



**Figure 2** A clinical pathway for the differential diagnosis of ear(s) involved and the differentiation of canalithiasis and cupulolithiasis for PC-BPPV.

or tapping of the mastoid process during the treatment procedure to facilitate movement of the debris. Our experience indicates that in patients requiring more than the average number of treatments, tapping the mastoid process has been somewhat successful in clearing the debris. A horizontal canal migration during, or subsequent to a Canalith Repositioning Maneuver, however, has occurred in greater numbers in those patients who have received the tapping.<sup>31</sup>

## POSTTREATMENT RECOMMENDATIONS

There are a wide variety of reported activity limitations for the patient from 24 to 48 hours following treatment. Many reports have asked that the patient not lie supine or tip their head for 48 hours following treatment. This is often assisted by wearing a soft cervical collar. Then, for several nights or up to a week following treatment, the patient is asked to refrain from lying on the treated or affected side while sleeping.

A modified prohibition of having the patient avoid tipping the head or lying supine for over 24 hours has proven to be sufficient in our clinic (Appendix C). This is supported by the work of Zucca,<sup>32</sup> who reported that otolith debris in the vestibular system of frogs would dissolve within the calcium-poor endolymph well

within the 24-hour period. Recent reports have suggested that it may not be necessary for any prohibition of head movement or activity.<sup>33</sup>

## COMPLICATIONS OF TREATMENT

Clinicians participating in the treatment and management of BPPV patients should recognize that several complications, although not common, could occur and when they do, they must be dealt with quickly and efficiently.

## HORIZONTAL CANAL MIGRATION

Migration of the otolith debris into the horizontal canal has been reported in the literature.<sup>34-41</sup> For audiologists performing ENG without the benefit of video-oculographic recordings, this phenomenon may be the explanation for a direction changing nystagmus seen in positional ENG sub tests.

The debris may migrate into the horizontal canal following Canalith Repositioning or during movements through the positioning. The patient will present with a burst of linear horizontal nystagmus with its fast phase beating geotropically towards the undermost involved horizontal canal. If the patient's head is turned onto the opposite or unaffected ear, there is a linear horizontal ageotropic nystagmus beating away from the normal undermost ear towards the affected ear. Patients may then

be treated with either the Appiani Liberatory, or a 360° barbeque roll maneuver. Appiani et al<sup>34</sup> described a simplified method of treatment for the horizontal canal BPPV migration. The treatment requires a side lying liberatory-lateral type movement followed by a downward head turn while the patient is in the lateral body position. This method is more comfortable and easily performed by heavier or older patients.

The barbeque roll technique requires the patient to roll in 90° quadrants until the debris is cleared. Both methods of treating the horizontal canal begin with the initial treatment position away from the affected ear. This differs from Canalith Repositioning and Semont Liberatory Maneuvers that initiate towards the affected ear.

### CANALITH JAM

A rare but often frightening occurrence is a canalith jam.<sup>44</sup> This occurs when the otolith debris is unable to clear the common crus as the debris falls downward from the posterior canal into the utricle when the patient returns to a seated position following the final stage of the Canalith Repositioning Maneuver or Semont Liberatory Maneuver. The patient will become symptomatic and the response will not be transient and it will not fatigue the patient. The sensation of falling or rapidly tumbling in a Tumarkin's-like crisis may be severe. The method used to clear the jam is to reverse the Repositioning protocol in the order in which it was performed.

### METHODS AND PATIENTS

A retrospective case review of 376 patients treated for PC-BPPV at the American Institute of Balance from May of 1994 to July of 2001 was conducted. To date, nearly 700 patients have been treated. The selection of the patients for this study was based on completeness of records and follow-up contact. All patients were treated by a single clinician. All patients had been medically referred for evaluation and treat-

ment of vertigo. Comprehensive cochleovestibular testing included ENG/VNG with air calorics. The diagnosis of PC-BPPV was made with the classic criteria of: (1) transient rotatory-torsional nystagmus toward the undermost ear, (2) subjective vertigo that parallels the nystagmus, (3) latency of onset of nystagmus, and (4) a possible reversal of nystagmus upon return to sitting position. Patients that were evaluated from 1994 to 1997 were confirmed by direct visual observation. Patients that were evaluated between 1998 and 2001 were video recorded with video-oculography (Micro Medical-monocular, SMI monocular video-goggles or SMI binocular 2D VOG). In instances where monocular recording was performed, the infrared camera was placed to record the ipsilateral eye of the undermost test ear. Figure 2 presents a clinical pathway for differential diagnosis of the ear(s) involved, and differentiation of canalithiasis vs. cupulolithiasis. Preceding all positioning maneuvers, a vertebral artery-screening test, as previously recommended by this author,<sup>30</sup> was performed. Positive indicators on a Vertebral Artery Screening test include: (1) diplopia, (2) dysarthria, (3) dizziness or near syncope, (4) nausea, and (5) in rare cases, nystagmus. A positive indicator on the Vertebral Artery Screening Test is reported to the referring physician, and the patient is not positioned with the neck hyperextended.

Modified Dix-Hallpike maneuvers in three variations were used to elicit the PC-BPPV response. The American Institute of Balance protocol excludes the traditional-classic Dix-Hallpike method due to the potentially negative biomechanical impact on both clinician and patient. The positioning techniques are: (1) modified Dix-Hallpike with hyperextension (head hanging with neck fully supported), (2) supine (without head hanging, fully supported) minimal hyperextension and head rotation, and (3) side-lying with minimal hyperextension and rotation. The positioning-test maneuver selection was based on patient comfort and on each patient's individual physiognomy or physical condition. A patient with a history of a fused vertebra, for example, is a poor candidate for any type of head hanging maneuver. Likewise, a patient with a recent hip



replacement (less than 90 days) is a poor side-lying candidate.

Patients identified as having HC-BPPV or AC-BPPV upon initial diagnosis or secondary to migration post initial treatment were excluded from the anecdotal review. All patients who were deemed candidates for treatment had medical clearance and/or a subsequent referral for treatment. Prior to treatment, patients received written materials describing their condition, reprints and articles detailing the treatment procedure, post treatment restrictions, possible complications, and the Institute's treatment outcomes. Treatment was never performed on the same day as the diagnostic evaluation.

## TREATMENT METHODS

Patients seen from 1994 to 1998 (N=272) were treated only with the Semont Liberatory Maneuver (SLM). Patients from 1999 to 2001 (N=195) were treated with the Canalith Repositioning Maneuver (CRM) as the first treatment of choice. The CRM patients who did not clear/fatigue on the initial treatment maneuver or those who were still positive for PC-BPPV on their first follow-up visit were treated with SLM. Descriptions of both treatment methods are presented in Appendices A and B for modified Canalith Repositioning Maneuver and Semont Liberatory Maneuver, respectively.

## POSTTREATMENT RESTRICTIONS AND FOLLOW-UP

Patients were provided with written instructions, as presented in Appendix C, restricting activity. Patients also were advised to return in approximately one week for a post treatment follow-up visit. Patients educated in advance of the importance of the follow-up visit(s) as part of their treatment were highly compliant. Their follow-up appointment was made at the time of their initial treatment and all visits were confirmed by telephone prior to their appointment day. A subjective report of treatment benefit by telephone inquiry has not

shown to be a reliable predictor of treatment efficacy. Patients often are reluctant to aggressively test themselves. While reporting that they are feeling better and have resumed normal activities, they may consciously or unconsciously avoid the most provocative head positions. Concomitant vestibular symptomatology (i.e., noncompensated high frequency vestibulopathy) or unrelated symptoms (i.e., lightheadedness) secondary to other medical conditions, may contaminate the patient's subjective report of treatment benefit.

Confirmation of extinction of the rotatory-torsional nystagmus and vertigo was obtained through the repetition of the diagnostic positioning maneuvers at the time of the patients' first follow-up visit. The patients were placed in at least two provocative positions. The first position was an appropriately modified Dix-Hallpike position, and the second was the side-lying position. Patients also were roll tested to ensure there had been no migration into the horizontal canal. A patient was considered cleared if no recordable nystagmus was noted through direct observation or video-oculography recording, and there was no subjective report of vertigo by the patient. If there was observable or recordable nystagmus and a subjective report of vertigo, even though significantly ameliorated, the patient was considered to remain positive for PC-BPPV, and re-treated.

## RESULTS

The 376 patients (Table 2) in the anecdotal review were comprised of 110 males (mean age 71 years), and 266 females (mean age 69 years). The patients received a total of 480 treatments. Of these, 195 received CRM, 272

**Table 2 Summary of 376 Patients by Gender, Age, and Involved Ear(s)**

	Age Range	Mean Age	Affected Ear		
			Right	Left	Both
Males	32-91	71	63	42	5
Females	28-91	69	148	106	12

**Table 3 Summary of 376 Patients' Outcomes by Treatment Method and Gender**

	Treatment			% Improved		
	Canalith Repositioning	Semont Liberatory	Both	Canalith Repositioning	Semont Liberatory	Both
Males	56	84	2	97.9	92.8	75
Females	139	188	11	98.9	96.8	95.9

received SLM, and 13 patients received both treatments (Table 3). The right ear was affected in 211 patients and the left ear in 148 patients. Both ears were affected in only 17 patients. There was a 56% predominance of a right ear involvement. The right ear predominance was not gender specific, with right ear involvement occurring in 57% of males, and in 56% of females.

As can be seen in Table 3, the treatment efficacy was comparable for males and females receiving both CRM and SLM treatments. The CRM treatment efficacy was approximately 98% in males, and 99% in females. SLM treatment efficacy was approximately 93% for males, and approximately 97% in females. For patients receiving both methods of treatment, there was a difference in the treatment efficacy, with females showing a positive outcome of approximately 96% versus males with 75%.

A consideration in the selection of treatment methods is the success rate with a single treatment. As indicated in Figure 6, 79% of the patients had a complete resolution following only one treatment (Table 4). Both males and females had a success rate of 98% with a single treatment using Canalith Repositioning maneuver. The Semont Liberatory Maneuver produced success in a single treatment in 91% of the males, and in approximately 97% of the females. Seventeen percent of the patients required two treatments, while 3.7% required three treatments. Only one-half of one percent

(0.5%) of the 376 patients required four treatments.

Another consideration of treatment method selection and follow-up is the recurrence rate. The data presented in Table 5 illustrate a recurrence rate for the Canalith Repositioning Maneuver of approximately 7% for both males and females. The Semont Liberatory Maneuver presented with a recurrence rate of only 3.6% and 1.5% for males and females, respectively. A recurrence was considered to have occurred 30 days following the initial treatment period and post follow-up period.

## DISCUSSION

A review of the literature over the past 14 years has shown the treatment efficacy of both Canalith Repositioning and Semont Liberatory Maneuvers. The patients in our anecdotal review with PC-BPPV demonstrated successful outcomes using both treatment methods. Seventy-nine percent of the patients were successfully cleared of symptoms with a single treatment and 17% required two treatments. These data compare favorably with Epley's finding that 80% of his 30 patients were successfully cleared with one treatment. Epley's data further suggested that 98% were symptom-free by the end of 3 months. Our data indicated that 96%

**Table 4 Comparison of Single Treatment Efficacy by Method and Gender**

	Canalith Repositioning	Semont Liberatory
Males	98.06	91.47
Females	98.41	96.78

**Table 5 Comparison of Recurrence Rate by Treatment Method and Gender**

	Recurrence Rate (%)		
	Canalith Repositioning	Semont Liberatory	Both
Males	7.2	3.6	0.9
Females	6.8	1.5	0

of the patients were successfully cleared following just two treatments. Semont et al reported similar findings with 84% of his patients requiring one treatment and 93% successfully cleared with two treatments.

Semont's recurrence rate over an 8-year period was approximately 4%. Our recurrence rates ranged between 2 to 4% using a Modified Semont Liberatory Maneuver, and 7% for the Canalith Repositioning Maneuver. To our knowledge, there has not been a longitudinal study that has reported a recurrence rate over an 8-year period with Canalith Repositioning Maneuver. Generally, there was no correlation between gender and treatment efficacy with any of the methods. The patient's mobility and compliance to the post-treatment restrictions are probably the two most important predictors of the patient's success. We did not differentiate the underlying pathophysiology or etiology of the BPPV. For example, would a patient with BPPV secondary to vestibular neuritis be less likely clear from a single treatment versus an idiopathic occurrence of BPPV? The high (79%) success rate of a single treatment, and a 96% success rate with just two treatments, suggests that it is unlikely that underlying etiology would provide information to affect treatment of choice.

## CONCLUSION

1. CRM and SLM are equally efficacious methods of treating PC-BPPV.
2. Ninety-six percent of patients will be successfully treated with two treatments. Seventy-nine percent require only a single treatment.
4. The recurrence rate of BPPV is relatively small (approximately 7%) using CRP, and as low as 2 to 4% using SLM.

Both CRM and SLM are highly efficacious treatment methods for PC-BPPV. Undiagnosed and untreated BPPV, particularly in the older population, is a significant and often catastrophic health hazard. The cost in dollars and human suffering in allowing patients to go untreated have been well documented.

With 96% of patients requiring two or less treatments, CRM and SLM are highly cost-effective methods in the treatment of this often-debilitating condition. Third-party payers, primary care physicians, and patients themselves should be aware of this highly efficacious and cost-effective treatment for the leading cause of dizziness and vertigo.

## ACKNOWLEDGMENTS

The authors wish to recognize The American Institute of Balance staff members Jeanne Berry and Denise Bednar for the manuscript preparation. University of South Florida doctoral students Mary Aguila, Erika Johnson, and Nancy Wong are recognized for their assistance in the data collection.

## ABBREVIATIONS

AC-BPPV	anterior canal benign paroxysmal positional vertigo
ASCC	anterior semicircular canal
BPPV	benign paroxysmal positional vertigo
CRM	Canalith Repositioning Maneuver
HC-BPPV	horizontal canal benign paroxysmal positional vertigo
HSCC	horizontal semicircular canal
PC-BPPV	posterior canal benign paroxysmal positional vertigo
PSCC	posterior semicircular canal
SLM	Semont Liberatory Maneuver
VNG	videonystagmography

## REFERENCES

1. Bath AP, Walsh RM, Ranalli P, et al. Experience from a multidisciplinary "dizzy" clinic. *Am J Otol* 2000;21:92-97
2. Li JC, Li CJ, Epley J, Weinberg L. Cost-effective management of benign positional vertigo using canalith repositioning. *Otolaryngol Head Neck Surg* 2000;122:334-339
3. Barany R. Diagnose von Krankheitserschernungen Bereiche des Otolithenapparaten. *Acta Otolaryngol (Stockh)* 1921;2:434-437
4. Dix MR, Hallpike CS. The pathology, symptomatology and diagnosis of certain common disorders

- of the vestibular system. *Ann Otol Rhinol Laryngol* 1952;61:987-1016
5. Schuknecht HF. Cupulolithiasis. *Arch Otolaryngol* 1969;90:113-126
  6. Baloh RW. Harold Schuknecht and pathology of the ear. *Otology and Neurotology* 2001;22:113-122
  7. Parnes LS, McClure JA. Free-floating endolymph particles: a new operative finding during posterior canal occlusion. *Laryngoscope* 1992;102: 988-992
  8. Hall SF, Rudy RRF, McClure JA. The mechanics of benign paroxysmal positional vertigo. *J Otolaryngol* 1979;8:151-158
  9. Gans RE. Overview of BPPV: pathophysiology and diagnosis. *The Hearing Review* 2000;7(8): 38-43
  10. Brandt T, Daroff RB. Physical Therapy for Benign Paroxysmal Positional Vertigo. *Arch Otolaryngol* 1980;106:484
  11. Semont A, Freyss G, Vitte E. Curing the BPPV with a Liberatory Maneuver. *Adv Otorhinolaryngol* 1988;42:390-393
  12. Epley JM. The Canalith Repositioning Procedure for treatment of benign paroxysmal positional vertigo. *Arch Otolaryngol* 1993;119:450-454
  13. Herdman SJ, Tusa RJ, Zee DS, et al. Single treatment approaches to benign paroxysmal positional vertigo. *Arch Otolaryngol* 1993;119:450-454
  14. Steenerson RL, Cronin GW. Comparison of the Canalith Repositioning Procedure and Vestibular Habituation Training in forty patients with benign paroxysmal positional vertigo. *Otolaryngol Head Neck Surg* 1996;114:61-64
  15. Coppo CF, Singarelli S, Fracchia P. Benign paroxysmal positional vertigo: follow-up of 165 cases treated by Semont's Liberating Maneuver. *Acta Otorhinolaryngol Ital* 1996;16:508-512
  16. Lempert T, Wolsley C, Davies R, Gresty MA, Bronstein AM. Three hundred sixty-degree rotation of the posterior semicircular canal for treatment of benign positional vertigo: a placebo-controlled trial. *Neurology* 1997;49:729-733
  17. Cohen HS, Jerabek J. Efficacy of treatments for posterior canal benign paroxysmal positional vertigo. *Laryngoscope* 1999;109:584-590
  18. Gall RM, Ireland DJ, Robertson DD. Subjective visual vertical in patients with benign paroxysmal positional vertigo. *J Otolaryngol* 1999;28:162-165
  19. Wolf JS, Boyev KP, Manockey DE. Success of the modified Epley Maneuver in treating benign paroxysmal positional vertigo. *Laryngoscope* 1999; 109:900-903
  20. Dornhoffer JL, Colvin GB. Benign paroxysmal positional vertigo and Canalith Repositioning: clinical correlations. *Am J Otol* 2000;21:230-233
  21. Macias JD, Lambert KM, Massingale S, Ellensohn A, Fritz JA. Variables affecting treatment in benign paroxysmal positional vertigo. *Laryngoscope* 2000; 110:1921-1924
  22. Nunez RA, Cass SP, Furman JM. Short and long term outcomes of Canalith Repositioning for benign paroxysmal positional vertigo. *J Otolaryngol* 2000;122:647-652
  23. O'Reilly R, Elford B, Slater R. Effectiveness of Particle Repositioning Maneuver in subtypes of benign paroxysmal positional vertigo. *Laryngoscope* 2000;110:1385-1388
  24. Tirelli G, D'Orlando E, Zarccone O, Giacomarro V, Russolo M. Modified Particle Repositioning Procedure. *Laryngoscope* 2000;100:462-467
  25. Oghalai JS, Manolidis S, Barth J, Stewart M, Jenkins HA. Unrecognized benign paroxysmal positional vertigo in elderly patients. *Otolaryngol Head Neck Surg* 2000;122;5:630-634
  26. Gans RE, Crandall C. Overview of BPPV: evaluating treatment outcomes with clinimetrics. *The Hearing Review* 2000;7(11):50-54
  27. Ware JE. How to Score the Revised MOS Short Form Health Scales (SF-36) Boston: The Health Institute, New England Medical Center Hospitals; 1988
  28. Herdman SJ, Tusa RJ. Complications of the Canalith Repositioning Procedure. *Arch Otolaryngol Head Neck Surg*, 1996;122,3:281-286
  29. Li JC. Mastoid Oscillation: a critical factor for success in Canalith Repositioning Procedure. *Otolaryngol Head Neck Surg* 1995;112:670-675
  30. Sargent EW, Bankaitis AE, Hollenbeak CS, Currens JW. Mastoid oscillation in canalith repositioning for paroxysmal positional vertigo. *Otology and Neurotology* 2001;22:205-209
  31. Gans RE. Overview of BPPV: treatment methodologies. *The Hearing Review* 2000;7(9):34-39
  32. Zucca, G, Valli S, Valli P, Perin P, Mira E. Why do benign paroxysmal positional vertigo episodes recover spontaneously? *J Vestib Res* 1998;8:325-329
  33. Nuti D, Nati C, Passali D. Treatment of benign paroxysmal positional vertigo: no need for post-manuever restrictions. *J Otolaryngol* 2000;122: 440-444
  34. Appiani GC, Catania G, Gagliardi M. A liberatory maneuver for the treatment of horizontal canal paroxysmal positional vertigo. *Otology and Neurotology* 2001;22:66-69
  35. Fife TD. Recognition and management of horizontal canal benign positional vertigo. *Am J Otol* 1998;19:345-351
  36. Vannucchi P, Giannoni B, Pagnini P. Treatment of horizontal semicircular canal benign paroxysmal positional vertigo. *J Vestib Res* 1997;7:1-6

37. DelaMeilleure G, Dehaene I, Depondt M, et al. Benign paroxysmal positional vertigo of the horizontal canal. *J Neurol Neurosurg Psychiatry* 1996; 60:68-71
38. Nuti D, Vannucchi P, Pagnini P. Benign paroxysmal positional vertigo of the horizontal canal: a form of canalolithiasis with variable clinical features. *J Vestib Res* 1996;6:173-184
39. Steddin S, Ing D, Brandt T. Horizontal canal benign paroxysmal positional vertigo (h-BPPV): transition of canalolithiasis to cupulolithiasis. *Ann Neurol* 1996;40:918-922
40. Strupp M, Brandt T, Steddin S. Horizontal canal benign paroxysmal positional vertigo: reversible ipsilateral caloric hypoexcitability caused by canalolithiasis. *Neurology* 1995;45:2072-2076
41. Lempert T, Tiel-Wick K. A positional maneuver for treatment of horizontal canal benign positional vertigo. *Laryngoscope* 1996;106:476-478
42. Baloh RW, Jacobson K, Honrubia V. Horizontal semicircular canal variant of benign positional vertigo. *Neurology* 1993;43:2542-2549
43. McClure JA. Horizontal canal BPPV. *J Otolaryngol* 1985;14:30-35
44. Herdman SJ, Tusa RJ. Complications of the Canalith Repositioning Procedure. *Arch Otolaryngol Head Neck Surg* 1996;122,3:281-286
45. Norre ME, Beckers A. Benign Paroxysmal Positional Vertigo in the elderly. *J Am Geriatr Soc* 1988;36:425-429
46. Troost BT, Patton JM. Exercise therapy for Positional Vertigo. *Neurology* 1992;42:1441-1444
47. Blatt PJ, Gergakakis GA, Herdman SJ, Clendaniel RA, Tusa RJ. The effect of the Canalith Repositioning Maneuver on resolving postural instability in patients with Benign Paroxysmal Positional Vertigo. *Am J Otol* 2000;21:356-363
48. Gross EM, Ress BD, Viirre ES, Nelson JR, Harris JP. Intractable Benign Paroxysmal Positional Vertigo in patients with Meniere's disease. *Laryngoscope* 2000;110:655-659

#### **APPENDIX A. DESCRIPTION OF MODIFIED CANALITH REPOSITIONING MANEUVER (CRM)**

**Step 1.** The patient is seated on the long axis of the examination table (we recommend placing a pillow under the legs to avoid cramping the hamstrings). The clinician stands at the head of the table to support the patient's neck and back. The patient turns the head *toward*

the affected ear as he or she lies backwards. The patient is then placed in the hyper-extended head hanging position while the clinician, sitting on a properly height-adjusted stool, provides full support for the patient's neck. Following the anticipated latency, there should be an onset of nystagmus and subjective vertigo. There must be a provocation in this initial position. The patient remains in this position for 3 minutes.

**Step 2.** The patient is then asked and cued to rotate the head to the opposite side while maintaining a hyper-extension with the clinician supporting the neck at all times. Regardless of whether there is a provocation, the patient maintains this position for 3 minutes.

**Step 3.** The patient now must roll onto his or her side (nonaffected ear side). This will place the patient face down. As in the hyper-extended, somewhat head-hanging position in Step 1, the head should be sufficiently off the table so the chin is extended beyond the edge of the table. The physics of this position is such that the head at this point no longer needs to be supported by the clinician. The position may cause a strong provocation, so it is imperative to provide sufficient kinesthetic feedback to the patient. The patient maintains this position for 3 minutes.

**Step 4.** The patient now swings the legs off the table and transfers to a fully seated, erect position by using elbow and hands to push themselves up. The head remains turned and is brought frontward only when the patient reaches the seated position. The head is then cued for a slight downward tilt and then returned to the normal or neutral position. The clinician must be prepared for the possibility that the patient may experience a Tumarkin's crisis. This is an intense sensation of falling or being pulled to the ground. The patient is placed in a soft cervical collar as a reminder to not move or pitch the head in the vertical plane.

## **APPENDIX B. DESCRIPTION OF MODIFIED SEMONT LIBERATORY MANEUVER TREATMENT**

**Step 1.** The patient is seated on the side of the examination table. The head is turned *away* from the affected ear and the side to be laid on.

**Step 2.** The patient is briskly laid on the side as they swing the legs up onto the table (similar movement as one does getting into bed). As the head has maintained the same position it is now resting on the table with the nose pointing upward at approximately a 45-degree angle. The patient maintains this position for five minutes following the cessation of nystagmus.

**Note:** The patient must be provoked (onset of nystagmus and vertigo) in the initial position. The nystagmus should appear following a brief latency. A few rare patients will show latencies as long as 50 seconds. The nystagmus and accompanying vertigo will last for 5 to 20 seconds.

**Step 3.** The patient is then moved to the opposite side through a transfer method whereby the legs hang from the table. The patient pushes up using elbow and hands, and is laid in the reverse direction on the opposite side. Because the patient has not moved the position of the head, the nose now faces down into the table. After waiting 10 seconds or so for a spontaneous burst of nystagmus and vertigo. If it does not occur, the patient is cued to move their head side to side. The head movement is initiated, whether there is nystagmus or not. The patient maintains this position for 5 minutes.

**Step 4.** The patient now swings the legs off the table and uses the elbows and hands to transfer to the seated-erect beginning position.

At this point, it is critical to maintain a physical hold of the patient for at least 20 seconds as, on occasion, patients may have a Tumarkins-like crisis. There is no way to predict in advance which patients may experience this intense sense of falling, dropping, or being pulled to the earth. The patient is then placed in a soft cervical collar for the sole purpose of serving as a friendly reminder to not move or pitch the head in the vertical plane.

## **APPENDIX C. POSTTREATMENT PATIENT INSTRUCTIONS**

### **Reminder on Day of Treatment**

1. Keep your head upright. Do not pitch your head up or down. Try to keep your head vertical, as if trying to balance a book on it.
2. Side-to-side turns are okay.

### **Sleeping for Tonight Only, and Next Day**

1. Do not lay flat. Try to sleep propped-up, about 30 degrees.
2. You may remove the cervical collar 24 hours post treatment.

### **Sleeping for the Next Three Days**

1. Avoid sleeping on the side of the treated ear.
2. Please call should you have any questions or concerns.
3. It is not unusual for you to feel a sensation of floaty-headedness, and you may be slightly off-balance for several days following treatment.

---

Patients are provided with a soft cervical collar.